

Antibiotic Susceptibility Profile of Klebsiella Species Isolated from Pregnant Women with Dysuria Attending Selected Medical Centre in Northern Nigeria

M. Umar^{1*}, I. A. Maidoki¹, A. A. Yaya², G. K. Ajiya³, S. P. E. Jakheng¹,
I. M. Abdulkarim⁴ and I. C. Amuta¹

¹Division of Microbiology, Department of Science Laboratory Technology, Nigerian Institute of Leather and Science Technology, PMB 1034, Zaria, Kaduna, Nigeria.

²Division of Biological Sciences, Department of Science Laboratory Technology, Nigerian Institute of Leather and Science Technology, PMB 1034, Zaria, Kaduna, Nigeria.

³Department of Research and Development, Nigerian Institute of Leather and Science Technology Extension Centre, Sokoto, Nigeria.

⁴Division of Biochemistry, Department of Science Laboratory Technology, Nigerian Institute of Leather and Science Technology, PMB 1034 Zaria, Kaduna, Nigeria.

Authors' contributions

This work was carried out in collaboration between all authors. Author MU designed the study, performed the statistical analysis, wrote the protocol, wrote the first draft of the manuscript. Authors IAM and AAY managed the analyses of the study. Authors SPEJ, IMA and ICA managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRRU/2018/43989

Editor(s):

(1) Dr. Dhastagir Sultan Sheriff, Professor, Department of Biochemistry, Faculty of Medicine, Benghazi University, Benghazi, Libya.

(2) Dr. Dragana Zivkovic, Professor, Department of Pediatric Urology, University of Novi Sad, Serbia.

Reviewers:

(1) Emmanuel Ifeanyi Obeagu, University Health Services, Michael Okpara University of Agriculture, Nigeria.

(2) Peter Daley, Memorial University, Canada.

(3) Maria Demetriou, Democritus University of Thrace, Greece.

Complete Peer review History: <http://www.sciencedomain.org/review-history/26568>

Original Research Article

Received 17th August 2018

Accepted 1st October 2018

Published 9th October 2018

ABSTRACT

Background: Urinary Tract Infection (UTI) is caused by the presence and growth of microorganisms in the urinary tract. Klebsiella infection is perhaps one of the common bacterial infections of mankind capable of causing urinary tract infections.

*Corresponding author: Email: mustamina19@gmail.com, mustapha4mina@yahoo.com;

Aims: This study targeted to evaluate the antibiotic susceptibility profile of *Klebsiella* species isolated from pregnant women reported with dysuria attending selected Medical Centre in Northern Nigeria.

Place and Duration of Study: The study was conducted at the Microbiology unit, Nigerian Institute of Leather and Science Technology, Zaria, and Ahmadu Bello University Health Service, Kaduna, Nigeria, between July 2010 and February 2011.

Methodology: Fifty mid-stream urine samples of were collected from consented pregnant women complaining of painful urination, attending selected medical centre in Northern Nigeria. *Klebsiella* species were isolated and biochemically characterized using standard microbiological methods. The isolates were subjected to antimicrobial sensitivity test using Kirby-Bauer disc diffusion technique.

Results: The prevalence of *Klebsiella* species was found to be 5 (10%). The *klebsiellae* species identified biochemically includes *Klebsiella oxytoca* and *K. pneumoniae*. The highest antimicrobial susceptibility in *Klebsiella* species was recorded by ciprofloxacin (100%) and gentamicin (100%). Resistance was recorded by ampicillin (0%). The antibiotics that were found to be more effective such as ciprofloxacin and gentamicin are regarded to be the first line drugs of choice for the treatment of urinary tract infections caused by the isolated *Klebsiella* species. Thus, continuous monitoring of antibacterial susceptibility before antibiotic prescription is important in order to monitor any emergence of resistance to the commonly active antimicrobials.

Conclusion: *Klebsiella oxytoca* and *K. pneumoniae* were isolated amongst the screened pregnant women with overall prevalence of 10%. From the results obtained, ciprofloxacin and gentamicin were found to be the first line drugs of choice for the treatment of urinary tract infections caused by the isolated *klebsiellae* species, whereas ampicillin recorded less activity.

Keywords: Antibacterial; biochemical characterizations; *Klebsiella*; urinary tract infection.

1. INTRODUCTION

Urinary Tract Infection (UTI) caused by the presence and growth of microorganisms in the urinary tract, is perhaps the single most common bacterial infection of mankind. Numerous reports have suggested that UTI can occur in both male and female patients of any age with bacterial counts of as low as 100 cfu/ml of urine [1].

Klebsiella is one of the common bacterial pathogens capable of causing urinary tract infections. The common pathogenic species of the genus *Klebsiella* includes: *Klebsiella pneumoniae*, *Klebsiella ozanae*, *Klebsiella oxytoca*, *Klebsiella rhinoscleromatis*, *Klebsiella terrigena*, *Klebsiella planticola* and *Klebsiella granulomatis* [2].

In female human subjects, urinary tract has an important association with the reproduction organ because of its proximity. In the non-pregnant state, the uterus lies just behind and partly over the bladder. While in the pregnant state, the enlarging uterus affects all the tissues of the urinary tract at various times. This is why urinary tract infection (UTI) is common in females. The highest incidence of urinary tract infection occurs in the child-bearing age and this has a direct relationship to sexual activity and aging [3]. UTI

may be asymptomatic in many cases, while it may be accompanied by dysuria (painful urination), cystitis (inflammation of urinary bladder) and pyelonephritis in other patients. The urethras and bladders normally prevent urine from backing up towards the kidneys and the flow of urine in the bladder helps wash bacteria out of the body. In men, the prostate gland produces secretions that slow bacterial growth. In both sexes, immune defenses also play a major role in the prevention of urinary infections. Despite these safeguards, and mechanisms, infection still occur [4]. Bacterial infections of the urinary tract are commonly seen in out-patients, hospitalized patients and apparently healthy populations [5]. A common source of infection is the catheters and tubes placed in the bladder. A person who cannot void, unconscious or critically ill, often need a catheter that stays in place for a long time. Some people, especially the elderly or those with nervous system disorders who lose bladder control, may need a catheter for life. Bacteria on the catheter can infect the bladder. So, it is imperative for the hospital staff to take special care to keep the catheter sterile and remove it as soon as possible [4]. In infants and children, the clinical presentation of UTI is often with non-specific clinical signs such as fever; irritability and vomiting that are also commonly seen in many acute self-limiting childhood viral

illness. UTI may be the signal of serious underlying congenital anomaly such as obstruction that may lead not only to more serious illness but also to renal damage. The infection may also be associated with progressive loss of kidney function either in association with renal dysphasia or with recurrent episodes of acute pyelonephritis [6].

Urinary Tract Infections caused by *Klebsiella* is one of the most devastating and fulminating disease that usually causes high rate of maternal mortality and morbidity in pregnant women, whom immunity is low, especially in rural areas of developing countries like Nigeria. Urinary tract infections are among the most common infections afflicting man. Urinary tract infections in young and adult women are usually uncomplicated, but are often recurrent and cause considerable morbidity. Urinary tract infection in pregnant women, elderly patients and catheterized patients warrant special attention because of their association with increased morbidity and possibly with increased mortality [7]. The antibiotic resistant strains of *Klebsiella* produce extended spectrum β -lactamase (ESBL) enzymes that destroy penicillin or cephalosporin classes of drugs, thereby conferring resistance to other commonly used antibiotic drug classes as well. These common bacteria, when they produce these enzymes are much harder to kill with antibiotics. The antibiotic resistance problem is likely to become widespread, as such it often become difficult to select an appropriate antibiotic therapy for urinary tract infection [8].

The aim of this study is to isolate, biochemically characterize and to determine the antibiotic susceptibility profile of *Klebsiella* species isolated from pregnant women with dysuria attending selected medical center in Northern Nigeria.

2. MATERIALS AND METHODS

Following reception of ethical approval from the Medical Advisory Committee of Ahmadu Bello University Health Service, Zaria, a total of 50 mid-stream urine samples were collected from consented pregnant women (previously confirmed to be pregnant by human chorionic gonadotrophin immunochromatographic test) suspected with various symptom as of urinary tract infections attending antenatal ward of Ahmadu Bello University Health Centre, Zaria. The urine samples were collected in screw-capped universal bottles and transported immediately to laboratory for analysis. Only

consented and volunteered pregnant women of various age groups suspected with symptoms of urinary tract infections such as cystitis or dysuria were enrolled in the study. Written consent forms were signed by the patients (Appendix-1). Non pregnant women, asymptomatic pregnant women and those symptomatic but not volunteered to participate in the study were excluded.

The portion of the urine samples were dispensed into sterile test-tubes and centrifuged at the speed of 1500 rpm for 15 minutes. The supernatant was discarded and the sediments were inoculated on the surface of prepared Cysteine Lactose Electrolyte Deficient (CLED) agar, then the suspected colonies of *Klebsiella* were sub-cultured on MacConkey agar and incubated at 37°C for 24 hours [9]. Following the incubation, *Klebsiella* species were isolated based on their macroscopic cultural characteristics (Plate 1). The suspected isolates were subjected to Gram's staining to determine the Gram potential and morphology of the isolated organisms. *Klebsiella* species were preliminarily identified based on their microscopic features as Gram negative bacilli (rod shaped), non-sporulating, and non-motile bacteria [10].

2.1 Biochemical Characterization of the Isolated *Klebsiella* Species

The following biochemical tests were employed in order to characterize the isolates up to their species level by comparing their reactions with that of the known taxa as documented in Bergey's manual of determinative keys in Bacteriology: Catalase, Methyl Red, Voges Proskauer, Indole, Citrate Utilization, motility, Oxidase, Urease, Nitrate Reduction, Esculin Hydrolysis and Triple Sugar Iron tests, as well as Lactose, Maltose, Sucrose and Glucose fermentations [10,11].

2.2 Antimicrobial Sensitivity Testing (Kirby-Bauer Disc Diffusion Method)

Kirby-Bauer disc diffusion method for antimicrobial sensitivity testing was employed. Muller Hinton agar was freshly prepared according to the manufacturer's specifications and dispensed in sterile petri-dishes and allowed to solidify. A 0.5 McFarland standard of the test organism suspension was aseptically inoculated onto the surface of Muller Hinton agar using sterile swabs. The commonly used antibiotic discs were mounted on the surface of the

inoculated plates with the aid of sterile forceps, and were allowed to diffuse for 5 minutes prior to incubation. The plates were incubated at 37°C for 24 hours. Antibiotics used include ampicillin (10 µg), erythromycin (15 µg), ciprofloxacin (5 µg), doxycycline (30 µg) and gentamicin (10 µg). Following the incubation, zones of growth inhibition were measured to the nearest millimeter (mm) and recorded as described by CLSI [12] and adopted by Umar et al. [13].

3. RESULTS AND DISCUSSION

Out the total 50 samples analyzed, a total number of 5 isolates were microscopically and biochemically identified as *Klebsiella* species. Other uropathogens were isolated as well (Table 1).

Table 2 shows the biochemical characterizations of the bacterial isolates obtained from pregnant women attending selected medical center in Northern Nigeria. The klebsiellae isolated include *Klebsiella oxytoca* and *K. pneumoniae*.

Table 3 and Fig. 1 show highest susceptibility of *Klebsiella* species to ciprofloxacin (100%) and gentamicin (100%) whereas the least susceptibility was recorded by ampicillin (0%). The antibiotics that were found to be more active against the test organisms are ciprofloxacin and gentamicin.

Urinary tract infections due to *Klebsiella* species recorded an overall prevalence of 10% (Table 1). This is relatively low prevalence compared to the studies of Oko et al. [14] and Aiyegoro [15] who reported a prevalence of 45.8% bacteriuria in pregnant women. *Klebsiella oxytoca* and *Klebsiella pneumoniae* were the organisms isolated from pregnant women attending Ahmadu Bello University Health Service. Even the prevalence recorded in this study may be due to the immune status of the pregnant women and some antenatal procedures that pregnant women often exposed to, such as frequent collection of high vaginal swabs and endocervical swabs by sharing unsterilized speculum during sample collection. This agrees with the findings of Martin et al. [16], Kevic [17] and Högenauer et al. [18] who reported *Klebsiella oxytoca* and *K. pneumoniae* as common bacterial species capable of causing nosocomial infections in immunocompromised individuals, especially during first trimester of pregnancy. This work

suggests that *Klebsiella* species is one of the aetiologic agents of urinary tract infection among the set of people screened during the time of this study.



Plate 1. Preliminary macroscopic morphology of *Klebsiella* species on McConkey agar

Table 1. Prevalence of the *Klebsiella* species from urinary tract of pregnant women attending selected medical centre in Northern Nigeria

Organism Isolated	Frequency of Isolates	Prevalence (%)
<i>Klebsiella species</i>	5	10
Other Isolates	23	46
No Growth	22	44
Total	50	

The preliminary microscopy, colonial morphology, coupled with biochemical characterizations (Table 2) were used for the isolation of the bacterial isolates up to their species level using Bergey's Manual for Bacteriology as guideline as described by Buchanan and Gibbons [11]. Gram negative non-motile rod shaped bacteria that showed positive reaction to catalase, Voges Proskauer, urease, citrate, nitrate reductase and esculin, which produce alkaline on the butt and slants of triple sugar iron medium, capable of fermenting lactose, maltose, sucrose and glucose were identified as *Klebsiella pneumoniae*. Similar bacterial isolates that are indole positive in addition to the features of *K. pneumoniae* were identified as *Klebsiella oxytoca*. All other isolates that did not conform to the aforementioned biochemical characteristics were identified as non klebsiellae isolates.

Table 2. Biochemical characterizations of the bacterial isolates obtained from pregnant women attending selected medical centre in Northern Nigeria

Microscopy	CAT	MR	VP	IND	CIT	MT	OXI	URE	NIT	ESC	TSI	Sugar fermentation				Inference
												LAC	MAL	SUC	GLU	
Gram-negative non sporing rods	+	-	+	-	+	-	-	+	+	+	A/A +Gas	+	+	+	+	<i>Klebsiella pneumoniae</i>
Gram-negative non sporing rods	+	-	+	+	+	-	-	+	+	+	A/A +Gas	+	+	+	+	<i>Klebsiella oxytoca</i>
Gram-negative rods	-	+	-	+	-	+	-	+	-	-	K/A+H ₂ S	-	-	+	+	Non-klebsiellae isolates

CAT= catalase; MR= methyl red; VP= Voges Proskauer; IND= indole; CIT= citrate; MT= motility; OXI= oxidase; URE= urease; NIT= nitrate reductase; ESC= esculin hydrolysis; TSI= triple sugar iron; Lac= lactose; MAL= maltose; SUC= sucrose; GLU= glucose; A= alkaline; K= acid; H₂S= hydrogen sulfide; + = positive; - = negative




Fig. 1. Antibiotic susceptibility of *Klebsiella* species

Table 3. Antibiotic susceptibility profile of *Klebsiella* species isolated from pregnant women attending selected medical centre in Northern Nigeria (n=5)

Antimicrobials	Disc potency (µg)	Susceptibility (%)	Resistance (%)
Ampicillin	10	0 (0)	5 (100)
Ciprofloxacin	5	5 (100)	0 (0)
Doxycycline	30	3 (60)	2 (40)
Erythromycin	15	1 (20)	4 (80)
Gentamicin	10	5 (100)	0 (0)

The antimicrobial susceptibility of the antibiotics used showed relatively higher level of susceptibility of *Klebsiella oxytoca* and *K. pneumoniae* to ciprofloxacin (100%) and gentamicin (100%) respectively (Table 3 and Fig. 1). This conformed to the work of Nwanze [19] and Awoniyi [20], who reported that *Klebsiella* isolates showed relatively higher level of susceptibility to ciprofloxacin and gentamicin. Antibiotic resistance observed in this study is a sufficient cause for serious concern, especially with resistance to ampicillin (0%). The resistance may be due to lateral acquisition of resistance plasmid. However, Martin et al. [16] reported that *Klebsiella oxytoca* is capable of acquiring antibiotic resistance, and the isolates have been shown to produce extended-spectrum beta-lactamases and carbapenemases that deactivate penicillin derivatives.

The high resistance showed by some antibiotics could be due to earlier exposure of the isolates to the drug which may have enhanced resistant development [20]. It could also be due to practices of self-medication and indiscriminate use of this antibiotics and the acquiring of plasmid encoded resistant genes [21]. The findings have no doubt highlighted the need for constant monitoring of susceptibility of specific pathogens in different populations of commonly used antimicrobial susceptibilities and to assist clinician in the rational choice of antibiotics therapy to prevent misused or over use of the antibiotics.

4. CONCLUSION

Klebsiella oxytoca and *K. pneumoniae* were isolated amongst the pregnant women with dysuria attending Ahmadu Bello University Health Service Clinic, with total prevalence of 10%. From the results obtained, ciprofloxacin and gentamicin can serve as drugs of choice for the management of UTI caused by the *Klebsiella* species, but ciprofloxacin was found to be generally more effective. While ampicillin recorded less activity against the test organisms.

Constant and frequent review of antibiotic susceptibility of *Klebsiella* should be carried out to check for possible emergence of drug resistance by the organism. There is need for continuous monitoring of bacterial antibiotic susceptibility before antibiotic prescription in order to ensure adequate treatment of urinary tract infection (UTI) caused by the bacteria, because blind treatment is not economically wise.

CONSENT

All authors declare that 'written informed consent was obtained from the patient (or other approved parties) for publication of this case report.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Theodore M. Prevalence and arteriogram of urinary tract infections among prison inmates in Nigeria. The Interact Journal of Microbiology. 2007;3(2):3-5. Umeh O, Leonard BB. *Klebsiella* Infections. Multi-drug interaction checker. Ann Intern Med. 2009;119(6): 454–460
- Inabo HI, Obanibi HBT. Antimicrobial susceptibility of some urinary tract clinical isolate to commonly used antibiotic. African Journal of Biotechnology. 2006; 5(5):487-489.

3. Barry M, Brenner Roger C, Wiggins Harry R, Jacobson David G, Warnock Ahmad Elbadawi, John T, Glayhack. Urinary tract infection in adults. National Kidney and Urologic Disease Information Clearing house; 1991.
4. Olaitan OJ. Asymptomatic bacteriuria in female student's population of a Nigeria University. The Interest Journal of Microbiology. 2006;2(2):1-4.
5. Andrew W. Urinary tract infection in children diagnosis, treatment and long term management. National Collaborating Centre for women's and Children's Health Clinical Guideline. 2007;20.
6. Hooton TM. The epidemiology of urinary tract infection and the concept of Significant Bacteriuria-Infection. 1990; 18(2):S40-S43.
7. Jorgensen J, Jan EP, James L, Monica H, Brian W. Drug-resistant *Escherichia coli* and *Klebsiella*. Bacteria in Hospital samples and elsewhere in US. Science daily; 2008.
8. Igwegbe AO, Ugboaja JO, Okoli OA. Prevalence of asymptomatic bacteriuria among pregnant women in Nnewi, southeast Nigeria. International Journal of Biological and Chemical Sciences. 2012; 6(1):88-98.
9. Cheesbrough M. District laboratory practice for tropical countries. Macmillan Publishing Company U.S.A. 2004;67-180.
10. Buchanan RE, Gibbons NE. Bergey's manual of determinative keys for bacteriology. 8th Edition. The Williams Co., Baltimore, USA. 1984;7-21.
11. CLSI. Clinical and laboratory Standards Institute. Performance standards for antimicrobial susceptibility testing; Twenty-Eighth Edition. Standards M02, M07 and M11. 950 West Valley Road, Suite 2500 Wayne, PA 19087 USA; 2018
12. Umar M, Arzai AH, Yusuf G, Oko JO, Jobbi DY. Serological characterization and antimicrobial sensitivity profile of *Haemophilus influenzae* serotypes isolated from Aminu Kano Teaching Hospital, Kano, Nigeria. British Microbiology Research Journal. 2016;15(5):1-10.
13. Oko JO, Abriba C, Umar M, Asitok AD, Audu JA, Jakheng SPE, Ojeleye FS, Amos AK. Antibiotics susceptibility study of uropathogens isolated from asymptomatic pregnant women attending a tertiary maternity clinic in Northern Nigeria. Journal of Complementary and Alternative Medical Research. 2017;2(3):1-9.
14. Aiyegoro OA, Igbinosa OO, Ogunmwonyi IN, Odjadlere CE, Igbinosa OE, Okoh AI. Incidence of urinary tract infection (UTI) among children and adolescents in Ile-Ife, Nigeria. Journal of Microbiology Research: 2007;013-019.
15. Martin H, Thomas V, Gernot Z, Benjamin W, Eva L, Helmut SJF, Josefa P, Robert K, Andrea GJ. Nosocomial outbreak of *Klebsiella pneumoniae* carbapenemase-producing *Klebsiella oxytoca* in Australia. Antimicrobial Agents and Chemotherapy. 2012;56(4):2158-2161. DOI: 10.1007/s00253-003-1444-y
16. Kevic S. Hospital infection problem persists, National Desk – 377; 2010.
17. Högenauer C, Langner C, Beubler E, et al. *Klebsiella oxytoca* as a causative organism of antibiotic-associated hemorrhagic colitis. N. Engl. J. Med., 2006;355(23):2418-2426. DOI: 10.1056/NEJMoa054765
18. Nwanze PI, Nwaro LM, Oranusi S, Demkpa U, Okwu MU, Babatunde BB, Anake TA, Jatto W, Asagwara CE. Urinary tract infection in Okada Village: Prevalence and antimicrobial susceptibility pattern. Scientific Research Essay. 2007;2(4):112 – 116.
19. Awoniyi DO, Oguntibeju OO. An epidemiological survey of neonatal sepsis in a hospital in Western Nigeria, African Journal of Microbiology Research. 2009;3 (6):385–389.
20. Akinyemi KO, Alabi SA, Taiwo A, Omonigbehin EA. Antimicrobial susceptibility pattern and plasmid profile of pathogenic bacteria isolated from subjects with urinary tract infection in Lagos Nigeria. Nigeria Quarterly Journal Hospital Medicine. 1997;1:7–11.

APPENDIX

**NIGERIAN INSTITUTE OF LEATHER AND SCIENCE TECHNOLOGY**
(NILEST)
Formerly, Federal College of Chemical and Leather Technology (CHELTECH)
(FEDERAL MINISTRY OF SCIENCE AND TECHNOLOGY)

DIRECTOR GENERAL/CEO:
ISUWA K. ADAMU B.Sc., M.Sc, Ph.D, MICCON, GAPA

Our Ref: NILEST/PER.532/VOL.I
Your Ref: _____

**P.M.B 1034,
ZARIA NIGERIA
E-mail: nilestzaria@yahoo.com
Website: www.nilest.org**

Date: 19th May, 2016

PATIENTS' CONSENT FORM

PATIENT/PATIENT'S PARENT OR GUARDIAN
I have been asked to give consent for myself/daughter/family member (strike out what is not applicable) to participate in this research study which will involve collection of urine sample in the completion of the research study. I have read the foregoing information, or it has been read to me. I have had the opportunity to ask questions about it and any questions that I have asked have been answered to my satisfaction. I consent voluntarily for myself/my child/my family member (strike out what is not applicable) to participate as a participant in this study. The results obtained thereby may be used in any way to improve the understanding and management of the urinary tract infections among pregnant women in our hospitals and community in general. Participation is strictly voluntary and refusal to participate in the research study will not in any way affect my right and benefit in this clinic/hospital.

Parent's age _____ Sex _____
Signature of Patient/Patient's guardian/Parent (if the patient is under 18 years) _____
Date _____

STATEMENT BY THE WITNESS
I have witnessed the accurate reading of the consent form to the families of the potential participant, and the individual has had the opportunity to ask questions. I confirm that the individual has given consent freely and individual data will be treated with strict confidentiality.

Name of witness _____ AND Thumb print of participant _____
Signature of witness _____
Date _____

STATEMENT BY THE RESEARCHER/PERSON TAKING CONSENT
I have accurately read out the information sheet to the patient or families of the potential participant, and to the best of my ability made sure that the person(s) understands that the following will be done:
1. Samples of urine will be collected using sterile container.
2. The samples will be analyzed in the laboratory to isolate *Klebsiella species*, and to determine the drugs for the treatment.
3. The findings may be documented for public enlightenment and medical intervention by the concerned authorities.

My names are Mustapha Umar (NILEST/PER/532), Division of Microbiology, Department Science Laboratory Technology, Zaria, Kaduna state, Nigeria. I am currently carrying out a research titled Antibiotic Susceptibility Profile of *Klebsiella* Species Isolated from Pregnant Women with Dysuria Attending Selected Medical Centre in Northern Nigeria. I confirm that the patient/parent/patient's guardian was given an opportunity to ask questions about the study, and all the questions asked by him/her have been answered correctly and to the best of my ability. I confirm that the individual has not been coerced into giving consent, and the consent has been given freely and voluntarily.

Name of Researcher/person taking the consent **UMAR, M.**

Appendix 1. Patients' informed consent form

© 2018 Umar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/26568>